This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.



Feeding organic containing wastewater into anaerobic bioreactor to let anaerobes therein decompose organic pollutants in wastewater;



Feeding effluent of anaerobic bioreactor into aerobic bioreactor where aerobes therein decompose residual organic in wastewater and carbon dioxide dissolved in water is stripped by air stripping effect that increases the pH value of water, whereby the crystals of metal carbonate, struvite and inorganic solids are produced and inserted in the floc matrix of aerobes through bioflocculation.



Feeding effluent of aerobic bioreactor into membrane separation reactor where solids in waster are separated to obtain effluent free of organic pollutants.




FIG. 3




FIG. 4




FIG. 5




FIG. 6